





## LARP4 rabbit pAb

Catalog No	YP-Ab-12309
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	LARP4 PP13296
Protein Name	LARP4
Immunogen	Synthesized peptide derived from human LARP4 AA range: 209-259
Specificity	This antibody detects endogenous levels of LARP4 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
0	1 mag/mal
Concentration	1 mg/ml
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment  sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment  sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Cytoplasm, Stress granule . Cytoplasm, cytosol . Localized throughout the cytosol. Partially localized in stress granules in response to arsenite treatment  sequence caution:Translation N-terminally extended.,similarity:Contains 1 HTH La-type RNA-binding domain.,similarity:Contains 1 RRM (RNA recognition motif) domain.,











## **Products Images**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).