

🕻 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

## DCNL3 rabbit pAb

Catalog No	YP-Ab-12401
Isotype	lgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	DCUN1D3
Protein Name	DCNL3
Immunogen	Synthesized peptide derived from human DCNL3 AA range: 205-255
Specificity	This antibody detects endogenous levels of DCNL3 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cell membrane . Cytoplasm . Nucleus . Cytoplasm, perinuclear region . After UVC treatment, the protein enters to the nucleus gradually (PubMed:18823379). Cell membrane localization is essential for CUL3 neddylation (PubMed:19617556)
Tissue Specificity	Tends to be down-regulated in different type of cancers, including lung neuroendocrine carcinoma, thyroid Huerthle cell carcinoma and lung squamous cell carcinoma (PubMed:25349211). Mostly expressed in testis and brain (PubMed:26906416). Highly expressed in liver, bladder and renal normal tissue than their tumor tissue counterparts (PubMed:18823379). Palmitoylation stabilizes DCUN1D3 at the cell membrane (PubMed:19617556).
Function	similarity:Contains 1 DCUN1 domain.,
Background	
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd


C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4°over night