



## DR5 rabbit pAb

<b>Catalog No</b>	YP-Ab-12414
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB; ELISA
<b>Gene Name</b>	TNFRSF10B DR5 KILLER TRAILR2 TRICK2 ZTNFR9 UNQ160/PRO186
<b>Protein Name</b>	DR5
<b>Immunogen</b>	Synthesized peptide derived from human DR5 AA range: 200-280
<b>Specificity</b>	This antibody detects endogenous levels of Human DR5
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Tumor necrosis factor receptor superfamily member 10B (Death receptor 5;TNF-related apoptosis-inducing ligand receptor 2;TRAIL receptor 2;TRAIL-R2;CD antigen CD262)
<b>Observed Band</b>	
<b>Cell Pathway</b>	Membrane; Single-pass type I membrane protein.
<b>Tissue Specificity</b>	Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain.
<b>Function</b>	disease:Defects in TNFRSF10B may be a cause of squamous cell carcinoma of the head and neck (HNSCC) [MIM:275355].,function:Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B.,induction:TNFRSF10B is regulated by the tumor suppressor p53.,similarity:Contains 1 death domain.,similarity:Contains 3 TNFR-Cys repeats.,subunit:Homotrimer. Can interact with TRADD and RIP.,tissue specificity:Widely expressed in adult and fetal tissues; very highly



expressed in tumor cell lines such as HeLa S3, K562, HL-60, SW480, A549 and G361; highly expressed in heart, peripheral blood lymphocytes, liv

**Background**

The protein encoded by this gene is a member of the TNF-receptor superfamily, and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL/APO-2L), and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene. [provided by RefSeq, Mar 2009],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images