





LAP1B Polyclonal Antibody

| Catalog No | YP-Ab-12739 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;ELISA;IHC |
| Gene Name | TOR1AIP1 |
| Protein Name | Torsin-1A-interacting protein 1 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human TOR1AIP1. AA range:431-480 |
| Specificity | LAP1B Polyclonal Antibody detects endogenous levels of LAP1B protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | TOR1AIP1; Torsin-1A-interacting protein 1; Lamin-associated protein 1B; LAP1B |
| Observed Band | 66kD |
| Cell Pathway | Nucleus inner membrane ; Single-pass membrane protein . |
| Tissue Specificity | Expressed in muscle, liver and kidney. |
| Function | function:Binds to A- and B-type lamins. Possible role in membrane attachment and assembly of the nuclear lamina.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the TOR1AIP family.,subunit:Interacts with TOR1A., |
| Background | This gene encodes a type 2 integral membrane protein that binds A- and B-type lamins. The encoded protein localizes to the inner nuclear membrane and may be involved in maintaining the attachment of the nuclear membrane to the nuclear lamina during cell division. Alternate splicing results in multiple transcript variants. |
| | [provided by RefSeq, Apr 2016], |
| matters needing attention | [provided by RefSeq, Apr 2016], Avoid repeated freezing and thawing! |








Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




19-



UpingBio technology Co.,Ltd

C Tel: 400-999-8863
■ Email:UpingBio@163.com





Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).