



Unc18-1 Polyclonal Antibody

Catalog No	YP-Ab-12835
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	STXBP1
Protein Name	Syntaxin-binding protein 1
Immunogen	The antiserum was produced against synthesized peptide derived from human MUNC-18a. AA range:279-328
Specificity	Unc18-1 Polyclonal Antibody detects endogenous levels of Unc18-1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	STXBP1; UNC18A; Syntaxin-binding protein 1; MUNC18-1; N-Sec1; Protein unc-18 homolog 1; Unc18-1; Protein unc-18 homolog A; Unc-18A; p67
Observed Band	
Cell Pathway	Cytoplasm, cytosol . Membrane; Peripheral membrane protein.
Tissue Specificity	Brain and spinal cord. Highly enriched in axons.
Function	disease:Defects in STXBP1 are the cause of early infantile epileptic encephalopathy type 4 (EIEE4) [MIM:612164]. Affected individuals have neonatal or infantile onset of seizures, suppression-burst pattern on EEG, profound mental retardation, and MRI evidence of hypomyelination.,function:May participate in the regulation of synaptic vesicle docking and fusion, possibly through interaction with GTP-binding proteins. Essential for neurotransmission and binds syntaxin, a component of the synaptic vesicle fusion machinery probably in a 1:1 ratio. Can interact with syntaxins 1, 2, and 3 but not syntaxin 4. May play a role in determining the specificity of intracellular fusion reactions.,similarity:Belongs to the STXBP/unc-18/SEC1 family.,subunit:Binds SYTL4 and STX1A.,tissue specificity:Brain and spinal cord. Highly enriched in axons.,
Background	This gene encodes a syntaxin-binding protein. The encoded protein appears to play a role in release of neurotransmitters via regulation of syntaxin, a



transmembrane attachment protein receptor. Mutations in this gene have been associated with infantile epileptic encephalopathy-4. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2010],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of NIH/3T3 cells, using MUNC-18a Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MUNC-18a Antibody. The picture on the right is blocked with the synthesized peptide.