



# mGluR2 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-13418
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	GRM2
<b>Protein Name</b>	Metabotropic glutamate receptor 2
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human GRM2. AA range:241-290
<b>Specificity</b>	mGluR2 Polyclonal Antibody detects endogenous levels of mGluR2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	GRM2; GPRC1B; MGLUR2; Metabotropic glutamate receptor 2; mGluR2
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cell membrane; Multi-pass membrane protein. Cell junction, synapse . Cell projection, dendrite .
<b>Tissue Specificity</b>	Detected in brain cortex (at protein level). Widely expressed in different regions of the adult brain as well as in fetal brain.
<b>Function</b>	function:Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity. May mediate suppression of neurotransmission or may be involved in synaptogenesis or synaptic stabilization.,similarity:Belongs to the G-protein coupled receptor 3 family.,subunit:Interacts with GRASP.,tissue specificity:Widely expressed in different regions of the adult brain as well as in fetal brain.,
<b>Background</b>	glutamate metabotropic receptor 2(GRM2) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5



and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Two transcript variants encoding different isoforms have been found for this gene


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunofluorescence analysis of LOVO cells, using GRM2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GRM2 Antibody. The picture on the right is blocked with the synthesized peptide.