

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

## Olfactory receptor 10Z1 Polyclonal Antibody

Catalog No	YP-Ab-13459
Isotype	lgG
Reactivity	Human;Monkey
Applications	WB;IF;ELISA
Gene Name	OR10Z1
Protein Name	Olfactory receptor 10Z1
Immunogen	The antiserum was produced against synthesized peptide derived from human OR10Z1. AA range:201-250
Specificity	Olfactory receptor 10Z1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10Z1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	OR10Z1; Olfactory receptor 10Z1; Olfactory receptor OR1-15
Observed Band	38kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	
Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Avoid repeated freezing and thawing!

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

