







Olfactory receptor 52E4 Polyclonal Antibody

OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors carising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor gene and proteins for this organism is independent of other organisms. [provided by		
Applications WB;IF;ELISA Gene Name OR52E4 Protein Name Olfactory receptor 52E4 Immunogen The antiserum was produced against synthesized peptide derived from huma OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor sing from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors ceptor genes and proteins for this organism is independent of other organisms, [provided by the offercent or the organisms, [provided by the offercent organisms, [provided by the offercent organisms, [provided by the organisms is dependent of other organisms, [provided by the organisms is dependent of other organisms, [provided by the organisms is dependent of other organ	Catalog No	YP-Ab-13569
Applications WB;IF;ELISA Gene Name OR52E4 Protein Name Olfactory receptor 52E4 Immunogen The antiserum was produced against synthesized peptide derived from huma OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptors enes and proteins for this organism is independent of other organisms, provided by Provided by	Isotype	IgG
Gene Name OR52E4 Protein Name Olfactory receptor 52E4 Immunogen The antiserum was produced against synthesized peptide derived from huma OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 profein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/2000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity-Belongs to the G-protein coupled receptor family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory recep	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 52E4 Immunogen The antiserum was produced against synthesized peptide derived from huma OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a remembers of a large family of G-protein-coupled receptors (GPCF arising from single coding-exon genes. Olfactory receptors gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Applications	WB;IF;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from huma OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCF arising from single coding-exon genes. Olfactory receptors gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Gene Name	OR52E4
OR52E4. AA range:200-249 Specificity Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors carising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor gene and proteins for this organism is independent of other organisms. [provided by	Protein Name	Olfactory receptor 52E4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors are members of a large family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [Provided by	Immunogen	The antiserum was produced against synthesized peptide derived from human OR52E4. AA range:200-249
Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor arising from single coding-exon genes. Olfactory receptors dare aneuronal response that triggers the perception of a smell. The olfactory receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein—mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Specificity	Olfactory receptor 52E4 Polyclonal Antibody detects endogenous levels of Olfactory receptor 52E4 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor .,similarity: Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Polyclonal, Rabbit,IgG
1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	
Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Synonyms OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55 Observed Band 35kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	OR52E4; Olfactory receptor 52E4; Olfactory receptor OR11-55
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	35kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory recept proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the large the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
RefSeq, Jul 2008],	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and










matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using Olfactory receptor 52E4 Polyclonal Antibody diluted at 1:2000



Immunofluorescence analysis of LOVO cells, using OR52E4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 and HeLa cells, using OR52E4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using OR52E4 antibody.