







## Olfactory receptor 5A1 Polyclonal Antibody

Observed Band  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		
Reactivity Human;Rat;Mouse;  Applications WB;ELISA  Gene Name OR5A1  Protein Name Olfactory receptor 5A1  Immunogen The antiserum was produced against synthesized peptide derived from human OR5A1. AA range:240-289  Specificity Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .;similarity:Belongs to the G-protein coupled receptor 1 family.,  Difactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms is independent of other organisms. [provided by proteins are in provided by proteins are assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Catalog No	YP-Ab-13580
Applications WB;ELISA  Gene Name OR5A1  Protein Name Olfactory receptor 5A1  Immunogen The antiserum was produced against synthesized peptide derived from human OR5A1. AA range:240-289  Specificity Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor 1 family.,  Background olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptor receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors roteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor semes and proteins for this organism is independent of other organisms, [provided by proteins for this organism is independent of other organisms, [provided by	Isotype	IgG
Gene Name         OR5A1           Protein Name         Olfactory receptor 5A1           Immunogen         The antiserum was produced against synthesized peptide derived from human OR5A1. AA range:240-289           Specificity         Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           Background         olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptor interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-ex	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 5A1  Immunogen The antiserum was produced against synthesized peptide derived from human OR5A1. AA range:240-289  Specificity Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a revenuenal response that triggers the perception of a smell. The olfactory receptors share a -7-transmembrane domain structure with many heurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Applications	WB;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human ORSA1. AA range:240-289 Specificity Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications. Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 5 subfamily A member 1 (OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a remembers of a large family of G-protein-coupled receptors (GPCR) arising from single coding-xon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransulter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Gene Name	OR5A1
Specificity Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor ,,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Protein Name	Olfactory receptor 5A1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms independent of other organisms. [provided by	Immunogen	
Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249           Observed Band         36kD           Cell Pathway         Cell membrane; Multi-pass membrane protein.           Tissue Specificity         Function           function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family.           Background         olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single codingers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Specificity	Olfactory receptor 5A1 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5A1 protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band  36kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organisms is independent of other organisms. [provided by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor 1 family.,  Background olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249         Observed Band       36kD         Cell Pathway       Cell membrane; Multi-pass membrane protein.         Tissue Specificity         Function       function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,         Background       olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band 36kD  Cell Pathway Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Storage Stability  -20°C/1 year  OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249  Observed Band  36kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Observed Band  36kD  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Observed Band  Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Multi-pass membrane protein.  Tissue Specificity  Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	OR5A1; OR5A1P; Olfactory receptor 5A1; OST181; Olfactory receptor OR11-249
Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Background  olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	36kD
Function  function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background  olfactory receptor family 5 subfamily A member 1(OR5A1) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	
	Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by





€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com




matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of MCF-7 cells using Olfactory receptor 5A1 Polyclonal Antibody



Western blot analysis of lysates from MCF-7 cells, using OR5A1 Antibody. The lane on the right is blocked with the synthesized peptide.