

Ø Website: www.upingBio.com

## CD70 Polyclonal Antibody

Catalog No	YP-Ab-13909
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CD70
Protein Name	CD70 antigen
Immunogen	The antiserum was produced against synthesized peptide derived from human CD70. AA range:68-117
Specificity	CD70 Polyclonal Antibody detects endogenous levels of CD70 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; CD27 ligand; CD27-L; Tumor necrosis factor ligand superfamily member 7; CD antigen CD70
Observed Band	30kD
Cell Pathway	Membrane ; Single-pass type II membrane protein .
Tissue Specificity	B-cell,Kidney,
Function	function:Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.,similarity:Belongs to the tumor necrosis factor family.,subunit:Homotrimer.,
Background	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis. [provided by RefSeq, Jul 2008],



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 💌 Email:UpingBio@163.com



## matters needing attention

Usage suggestions

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

