



# CaMKV Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-14692
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	CAMKV
<b>Protein Name</b>	CaM kinase-like vesicle-associated protein
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CAMK5. AA range:211-260
<b>Specificity</b>	CaMKV Polyclonal Antibody detects endogenous levels of CaMKV protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CAMKV; CaM kinase-like vesicle-associated protein
<b>Observed Band</b>	54kD
<b>Cell Pathway</b>	Cell membrane ; Peripheral membrane protein . Cytoplasmic vesicle membrane ; Peripheral membrane protein . Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be associated with vesicular structures (By similarity) .
<b>Tissue Specificity</b>	Brain,Lung,Retinoblastoma,Teratocarcinoma,
<b>Function</b>	cofactor:Calcium.,domain:The protein kinase domain is predicted to be catalytically inactive.,function:Does not appear to have detectable kinase activity.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,subcellular location:Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be associated with vesicular structures.,subunit:Interacts with calmodulin, in the presence of calcium.,
<b>Background</b>	cofactor:Calcium.,domain:The protein kinase domain is predicted to be catalytically inactive.,function:Does not appear to have detectable kinase activity.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,subcellular



location:Predominantly observed in association with the plasma membrane of soma and in neurites, both axons and dendrites. May be associated with vesicular structures.,subunit:Interacts with calmodulin, in the presence of calcium.,


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


**Products Images**




Western Blot analysis of various cells using CaMKV Polyclonal Antibody



Immunofluorescence analysis of HeLa cells, using CAMK5 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using CAMK5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 and HUVEC cells, using CAMK5 Antibody. The lane on the right is blocked with the synthesized peptide.