

Website: www.upingBio.com

MAST3 Polyclonal Antibody

Catalog No YP-Ab-14828 Isotype IgG Reactivity Human;Mouse Applications WB;ELISA Gene Name MAST3 Protein Name Microtubule-associated serine/threonine-protein kinase 3 Immunogen The antiserum was produced against synthesized peptide derived from h MAST3. AA range:1031-1080 Specificity MAST3 Polyclonal Antibody detects endogenous levels of MAST3 proteir Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit.IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase Observed Band 143kD Cell Pathway intracellular, Tissue Specificity Brain, Function catalytic activity.ATP + a protein = ADP + a phosphoprotein., caution:It is u whether Met-1/Met-172 is the initiatorordactor/Magnesium, similarity.	
Reactivity Human;Mouse Applications WB;ELISA Gene Name MAST3 Protein Name Microtubule-associated serine/threonine-protein kinase 3 Immunogen The antiserum was produced against synthesized peptide derived from h MAST3 AA range: 1031-1080 Specificity MAST3 Polyclonal Antibody detects endogenous levels of MAST3 protein Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications: Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase Observed Band 143kD Cell Pathway intracellular, Tissue Specificity Brain, Function catalytic activity:XTP + a protein = ADP + a phosphoprotein, caution:1; is travellular, ismilarity:Contains 1 PD2 (OHR) domain, similarity:Contains 1 protein kinase sup AGC Ser/Thr protein kinase family, similarity:Contains 1 protein kinase sup AGC Ser/Thr protein kinase family, si	
Applications WB;ELISA Gene Name MAST3 Protein Name Microtubule-associated serine/threonine-protein kinase 3 Immunogen The antiserum was produced against synthesized peptide derived from h MAST3. AA range:1031-1080 Specificity MAST3 Polyclonal Antibody detects endogenous levels of MAST3 protein Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase Observed Band 143kD Cell Pathway Brain, Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, caution:11 is u whether Met-17, Met-171 or Met-172 is the initiator, cofactor.Magnesium, similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family, similarity:Contains 1 AGC-kinase C-de domain, similarity:Contains 1 PD2 (DHR) domain, similarity:Contains 1 protein kinase domain, subunit.1Interacts with PTEN.,	
Gene NameMAST3Protein NameMicrotubule-associated serine/threonine-protein kinase 3ImmunogenThe antiserum was produced against synthesized peptide derived from h MAST3. AA range:1031-1080SpecificityMAST3 Polyclonal Antibody detects endogenous levels of MAST3 protein FormulationFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid SourcePurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase whether Met-1, Met-171 or Met-172 is the initiator, cortactor/magnesium, similarity:Belongs to the protein kinase sup AGC Ser/Th protein #ADP + a phosphoprotein., caution:11 is u whether Met-1, Met-171 or Met-172 is the omilarity.contains 1 AGC-kinase C-ed domain., similarity.contains 1 AGC-kinase C-ed domain., simi	
Protein Name Microtubule-associated serine/threonine-protein kinase 3 Immunogen The antiserum was produced against synthesized peptide derived from he MAST3. AA range:1031-1080 Specificity MAST3 Polyclonal Antibody detects endogenous levels of MAST3 protein Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinae Observed Band 143kD Cell Pathway intracellular, Tissue Specificity Brain, Function catalytic activity:ATP + a protein = ADP + a phosphoproteincaution:It is u whether Met-1, Met-171 or Met-172 is the protein kinase sup AGC Ser/Thr protein kinase familysimilarity:Contains 1 AGC-kinase C-te domainsimilarity:Contains 1 AGC-kinase C-te domainsimilarity:Contains 1 PDZ (DHR) domainsimilarity:Contains 1 AGC-kinase C-te domainsimilarity:Contains 1 PDZ (DHR) domainsimilarity:Contains 1 AGC	
ImmunogenThe antiserum was produced against synthesized peptide derived from h MAST3. AA range:1031-1080SpecificityMAST3 Polyclonal Antibody detects endogenous levels of MAST3 protein FormulationEiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid SourcePolyclonal, Rabbit, IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinae intracellular,Discus SpecificityBrain,FunctionCetalytic activity:ATP + a protein = ADP + a phosphoprotein., caution:It is u winitiator., cofactor:Magnesium., similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family., similarity:Contains 1 AGC-kinase C-te domain., similarity:Contains 1 PDZ (DHR) domain., similarity:Contains 1 AGC-kinase C-te domain., similarity:Contains 1 PDZ (DHR)	
MAST3. AA range:1031-1080SpecificityMAST3 Polyclonal Antibody detects endogenous levels of MAST3 proteinFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azidSourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase Observed BandObserved Band143kDCell PathwayBrain,Functiongrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein., caution:It is u wintratorofactor:Magnesium., similarity:Contains 1 AGC-kinase C-te domain., similarity:Contains 1 AGC-kinase C-te	
FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azidSourcePolyclonal, Rabbit, IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinasObserved Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-17 lor Met-172 is the initiatry.contains 1 PD2 (DHR) domain.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PD2 (DHR) domain.,similarity:Contains 1 pr	numan
SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinaeObserved Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein., caution:It is u whether Met-1, Met-172 is the initiator., cofactor:Magnesium., similarity:Selongs to the protein kinase sup AGC Ser/Thr protein kinase family., similarity:Contains 1 pd/Contains 1 pd/	in.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinas: Observed Band 143kD Cell Pathway intracellular, Tissue Specificity Brain, Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 PDZ (DHR)	ide.
affinity-chromatography using epitope-specific immunogen.DilutionWestern Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinase Observed BandObserved Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Contains 1 AGC-kinase c-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 PDZ	
applications.Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinae:Observed Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
Purity ≥90% Storage Stability -20°C/1 year Synonyms MAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinast Observed Band 143kD Cell Pathway intracellular, Tissue Specificity Brain, Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,coactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
Storage Stability-20°C/1 yearSynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinaseObserved Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
SynonymsMAST3; KIAA0561; Microtubule-associated serine/threonine-protein kinasObserved Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
Observed Band143kDCell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr	
Cell Pathwayintracellular,Tissue SpecificityBrain,Functioncatalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	ase 3
Tissue Specificity Brain, Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,caution:It is u whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	
Background catalytic activity: $\Delta TP + a$ protein = $\Delta DP + a$ phosphoprotein, caution: It is i	perfamily rerminal
whether Met-1, Met-171 or Met-172 is the initiator.,cofactor:Magnesium.,similarity:Belongs to the protein kinase sup AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-te domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 pr kinase domain.,subunit:Interacts with PTEN.,	



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com



matters needing attention

Usage suggestions

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

