



PI 3-Kinase p110 γ Polyclonal Antibody

Catalog No	YP-Ab-14914
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;IHC;IF;ELISA
Gene Name	PIK3CG
Protein Name	Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit gamma isoform
Immunogen	The antiserum was produced against synthesized peptide derived from human PIK3CG. AA range:881-930
Specificity	PI 3-Kinase p110 γ Polyclonal Antibody detects endogenous levels of PI 3-Kinase p110 γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PIK3CG; Phosphatidylinositol 4; 5-bisphosphate 3-kinase catalytic subunit gamma isoform; PI3-kinase subunit gamma; PI3K-gamma; PI3Kgamma; PtdIns-3-kinase subunit gamma; Phosphatidylinositol 4,5-bisphosphate 3-kinase 110 kDa catalytic subunit
Observed Band	120kD
Cell Pathway	Cytoplasm . Cell membrane .
Tissue Specificity	Pancreas, skeletal muscle, liver and heart.
Function	catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate. ,enzyme regulation:Activated by both the alpha and the beta-gamma G proteins.,function:3-phosphorylates the cellular phosphoinositide PtdIns-4,5-biphosphate (PtdIns(4,5)P2) to produce PtdIns-3, 4,5-triiphosphate (PtdIns(3,4,5)P3). Links G-protein coupled receptor activation to the secondary messenger PtdIns(3,4,5)P3 production.,pathway:Phospholipid metabolism; phosphatidylinositol phosphate biosynthesis.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Heterodimer of a catalytic subunit (PIK3CG/p120) and a regulatory (PIK3R5a/p101) subunit.,tissue specificity:Pancreas, skeletal muscle, liver and heart.,



Background

Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha, p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2015],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Western blot analysis of PIK3CG Antibody. The lane on the right is blocked with the PIK3CG peptide.



Immunohistochemistry analysis of paraffin-embedded human pancreas, using PIK3CG Antibody. The lane on the right is blocked with the PIK3CG peptide.



Western blot analysis of the lysates from Jurkat cells using PIK3CG antibody.