



Actin, Muscle Specific (ABT-MSA) mouse mAb

Catalog No	YP-Ab-15233
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;WB;IF
Gene Name	Actin, Muscle Specific
Protein Name	Actin, Muscle Specific
Immunogen	Synthesized peptide derived from human Actin, Muscle Specific
Specificity	This antibody detects endogenous levels of human Actin, Muscle Specific. Heat-induced epitope retrieval (HIER) Citrate buffer of pH6.0 was highly recommended as antigen repair method in paraffin secti
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Mouse, Monoclonal/IgG1, Kappa
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:100-500, WB 1:200-1000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm, cytoskeleton.
Tissue Specificity	Muscle,Tongue,
Function	disease:Defects in ACTC1 are the cause of cardiomyopathy dilated type 1R (CMD1R) [MIM:102540]. Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature death.,disease:Defects in ACTC1 are the cause of cardiomyopathy familial hypertrophic type 11 (CMH11) [MIM:612098]. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.,function:Actins are highly conserv
Background	Actins are highly conserved proteins that are involved in various types of cell motility. Polymerization of globular actin (G-actin) leads to a structural filament



(F-actin) in the form of a two-stranded helix. Each actin can bind to four others. The protein encoded by this gene belongs to the actin family which is comprised of three main groups of actin isoforms, alpha, beta, and gamma. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. Defects in this gene have been associated with idiopathic dilated cardiomyopathy (IDC) and familial hypertrophic cardiomyopathy (FHC). [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Human appendix tissue was stained with anti-MSA(ABT-MSA) antibody. Secondary Antibody was Goat anti Rabbit/Mouse polymer HRP, Ready to Use(RS0011) at 37° 45min.




Human cardiac muscle tissue was stained with anti-MSA(ABT-MSA) antibody. Secondary Antibody was Goat anti Rabbit/Mouse polymer HRP, Ready to Use(RS0011) at 37° 45min.



Human leiomyoma tissue was stained with anti-MSA(ABT-MSA) antibody. Secondary Antibody was Goat anti Rabbit/Mouse polymer HRP, Ready to Use(RS0011) at 37° 45min.



Human rhabdomyosarcoma tissue was stained with anti-MSA(ABT-MSA) antibody. Secondary Antibody was Goat anti Rabbit/Mouse polymer HRP, Ready to Use(RS0011) at 37° 45min.



Human skeletal muscle tissue was stained with anti-MSA(ABT-MSA) antibody. Secondary Antibody was Goat anti Rabbit/Mouse polymer HRP, Ready to Use(RS0011) at 37° 45min.