



# Topoisomerase II $\alpha$ mouse mAb(ABT272)

<b>Catalog No</b>	YP-Ab-15627
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	IHC;WB;IF
<b>Gene Name</b>	TOP2A TOP2
<b>Protein Name</b>	Topoisomerase II $\alpha$
<b>Immunogen</b>	Synthesized peptide derived from human Topoisomerase II $\alpha$
<b>Specificity</b>	The antibody can specifically recognize human Topoisomerase II $\alpha$ protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.82% sodium azide.
<b>Source</b>	Mouse, Monoclonal/IgG1, Kappa
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
<b>Dilution</b>	IHC-p 1:100-500, WB 1:200-1000, IF 1:100-500
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	$\geq 90\%$
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	DNA topoisomerase 2-alpha (EC 5.99.1.3;DNA topoisomerase II, alpha isozyme)
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Nucleus, nucleoplasm . Nucleus . Nucleus, nucleolus .
<b>Tissue Specificity</b>	Expressed in the tonsil, spleen, lymph node, thymus, skin, pancreas, testis, colon, kidney, liver, brain and lung (PubMed:9155056). Also found in high-grade lymphomas, squamous cell lung tumors and seminomas (PubMed:9155056).
<b>Function</b>	catalytic activity:ATP-dependent breakage, passage and rejoining of double-stranded DNA.,enzyme regulation:Specifically inhibited by the intercalating agent amsacrine.,function:Control of topological states of DNA by transient breakage and subsequent rejoining of DNA strands. Topoisomerase II makes double-strand breaks.,miscellaneous:Eukaryotic topoisomerase I and II can relax both negative and positive supercoils, whereas prokaryotic enzymes relax only negative supercoils.,PTM:Phosphorylation has no effect on catalytic activity.,similarity:Belongs to the type II topoisomerase family.,subcellular location:Generally located in the nucleoplasm.,subunit:Homodimer. Interacts with COPS5.,
<b>Background</b>	This gene encodes a DNA topoisomerase, an enzyme that controls and alters the topologic states of DNA during transcription. This nuclear enzyme is involved in processes such as chromosome condensation, chromatid separation, and the relief of torsional stress that occurs during DNA transcription and replication. It



catalyzes the transient breaking and rejoining of two strands of duplex DNA which allows the strands to pass through one another, thus altering the topology of DNA. Two forms of this enzyme exist as likely products of a gene duplication event. The gene encoding this form, alpha, is localized to chromosome 17 and the beta gene is localized to chromosome 3. The gene encoding this enzyme functions as the target for several anticancer agents and a variety of mutations in this gene have been associated with the development of drug resistance. Reduced activity of this enzyme may also pla

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images




Human lung squamous cell carcinoma tissue was stained with Anti-Topoisomerase II $\alpha$  (ABT272) Antibody




Human lymphoma tissue was stained with Anti-Topoisomerase II $\alpha$  (ABT272) Antibody



Human seminoma tissue was stained with Anti-Topoisomerase II $\alpha$  (ABT272) Antibody



Human testis tissue was stained with Anti-Topoisomerase II $\alpha$  (ABT272) Antibody



Human tonsil tissue was stained with  
Anti-Topoisomerase II $\alpha$  (ABT272) Antibody