

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



HoxD8 Polyclonal Antibody

Catalog No	YP-Ab-15771
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	HOXD8
Protein Name	Homeobox protein Hox-D8
Immunogen	The antiserum was produced against synthesized peptide derived from human HOXD8. AA range:241-290
Specificity	HoxD8 Polyclonal Antibody detects endogenous levels of HoxD8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HOXD8; HOX4E; Homeobox protein Hox-D8; Homeobox protein Hox-4E; Homeobox protein Hox-5.4
Observed Band	32kD
Cell Pathway	Nucleus.
Tissue Specificity	Kidney,Peripheral Nervous System,
Function	function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.,similarity:Belongs to the Antp homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,
Background	This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conserved family of transcription factors that play an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, located on different chromosomes, consisting of 9 to 11 genes arranged in tandem. This gene is one of several homeobox HOXD genes located in a cluster on chromosome 2. Deletions that remove the entire HOXD gene cluster or the 5' end of this cluster have been associated with severe limb and genital abnormalities. In addition to effects during embryogenesis, this particular gene



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



may also play a role in adult urogenital tract function. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Dec 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of HOXD8 Antibody. The lane on the right is blocked with the HOXD8 peptide.