

# **BMP-5** Polyclonal Antibody

YP-Ab-15879
lgG
Human;Mouse
WB;IHC;IF;ELISA
BMP5
Bone morphogenetic protein 5
The antiserum was produced against synthesized peptide derived from human BMP-5. AA range:114-163
BMP-5 Polyclonal Antibody detects endogenous levels of BMP-5 protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200
1 mg/ml
≥90%
-20°C/1 year
BMP5; Bone morphogenetic protein 5; BMP-5
52kD
Secreted.
Expressed in the lung and liver.
function:Induces cartilage and bone formation.,online information:Bone morphogenetic protein 5 entry,similarity:Belongs to the TGF-beta family.,subunit:Homodimer; disulfide-linked.,tissue specificity:Expressed in the lung and liver.,
This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone and cartilage development. Polymorphisms in this gene may be associated with osteoarthritis in human patients. This gene is differentially regulated in multiple human cancers. This gene encodes distinct protein isoforms that may be similarly proteolytically processed. [provided by RefSeq, Jul 2016],



### UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com




matters needing attention

Usage suggestions


#### Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using BMP-5 Polyclonal Antibody diluted at 1:1000






## UpingBio technology Co.,Ltd

🕲 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

WebsIte: www.upingBio.com



	-117 -85
BMP-5	-49
	-34
	-25
	-25

Western blot analysis of lysate from HUVEC cells, using BMP-5 antibody.

Immunohistochemistry analysis of BMP-5 antibody in

paraffin-embedded human heart tissue.