

CCK Polyclonal Antibody

Catalog No	YP-Ab-15885
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	ССК
Protein Name	Cholecystokinin
Immunogen	The antiserum was produced against synthesized peptide derived from human CCK. AA range:46-95
Specificity	CCK Polyclonal Antibody detects endogenous levels of CCK protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CCK; Cholecystokinin; CCK
Observed Band	
Cell Pathway	Secreted.
Tissue Specificity	Brain,Pancreas,
Function	function: This peptide hormone induces gall bladder contraction and the release of pancreatic enzymes in the gut. Its function in the brain is not clear. Binding to CCK-A receptors stimulates amylase release from the pancreas, binding to CCK-B receptors stimulates gastric acid secretion.,online information: Cholecystokinin entry, PTM: The precursor is cleaved by proteases to produce a number of active cholecystokinins.,similarity: Belongs to the gastrin/cholecystokinin family.,subunit: Binds to CCK-A receptors in the pancreas and CCK-B receptors in the brain.,
Background	This gene encodes a member of the gastrin/cholecystokinin family of proteins. The encoded preproprotein is proteolytically processed to generate multiple protein products, including the peptide hormones cholecystokinin-8, -12, -33, and others. The encoded peptides have been shown to regulate gastric acid secretion and food intake. A sulfated form of cholecystokinin-8 may modulate neuronal activity in the brain. Alternative splicing results in multiple transcript variants.




UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

[provided by RefSeq, Nov 2015],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of CCK antibody in paraffin-embedded human brain tissue.

Website: www.upingBio.com