



VIP Polyclonal Antibody

Catalog No	YP-Ab-15982
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;WB;ELISA
Gene Name	VIP
Protein Name	VIP peptides
Immunogen	The antiserum was produced against synthesized peptide derived from human VIP. AA range:71-120
Specificity	VIP Polyclonal Antibody detects endogenous levels of VIP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	VIP; VIP peptides
Observed Band	20kD
Cell Pathway	Secreted.
Tissue Specificity	Pancreatic carcinoma,Pheochromocytoma,Prostate,
Function	function:PHM and PHV also cause vasodilation.,function:VIP causes vasodilation, lowers arterial blood pressure, stimulates myocardial contractility, increases glycogenolysis and relaxes the smooth muscle of trachea, stomach and gall bladder.,online information:Vasoactive intestinal peptide entry,similarity:Belongs to the glucagon family.,
Background	The protein encoded by this gene belongs to the glucagon family. It stimulates myocardial contractility, causes vasodilation, increases glycogenolysis, lowers arterial blood pressure and relaxes the smooth muscle of trachea, stomach and gall bladder. The protein also acts as an antimicrobial peptide with antibacterial and antifungal activity. Alternative splicing occurs at this locus and two transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Nov 2014],


**matters needing attention**

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images



Immunohistochemical analysis of paraffin-embedded Human testis. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Immunohistochemistry analysis of VIP antibody in paraffin-embedded human testis tissue.



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night