



# CLIC6 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-16406
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	CLIC6
<b>Protein Name</b>	Chloride intracellular channel protein 6
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CLIC6. AA range:411-460
<b>Specificity</b>	CLIC6 Polyclonal Antibody detects endogenous levels of CLIC6 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CLIC6; CLIC1L; Chloride intracellular channel protein 6; Parchorin
<b>Observed Band</b>	75kD
<b>Cell Pathway</b>	Cytoplasm . Cell membrane ; Single-pass membrane protein . Predominantly cytoplasmic. Upon chloride ion efflux from the cell, it is translocated to the plasma membrane (By similarity). .
<b>Tissue Specificity</b>	Expressed in brain, placenta, pancreas and liver.
<b>Function</b>	domain:Members of this family may change from a globular, soluble state to a state where the N-terminal domain is inserted into the membrane and functions as chloride channel. A conformation change of the N-terminal domain is thought to expose hydrophobic surfaces that trigger membrane insertion.,function:May insert into membranes and form chloride ion channels. May play a critical role in water-secreting cells, possibly through the regulation of chloride ion transport.,PTM:Phosphorylated.,similarity:Belongs to the chloride channel CLIC family.,similarity:Contains 1 GST C-terminal domain.,subcellular location:Predominantly cytoplasmic. Upon chloride ion efflux from the cell, it is translocated to the plasma membrane.,subunit:Interacts with dopamine receptors DRD2, DRD3 and DRD4.,tissue specificity:Expressed in brain, placenta, pancreas and liver.,

**Background**

chloride intracellular channel 6(CLIC6) Homo sapiens This gene encodes a member of the chloride intracellular channel family of proteins. The gene is part of a large triplicated region found on chromosomes 1, 6, and 21. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Nov 2015],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of CLIC6 Antibody. The lane on the right is blocked with the CLIC6 peptide.



Western blot analysis of the lysates from HepG2 cells using CLIC6 antibody.