

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com



## HMGCL rabbit pAb

| Catalog No                | YP-Ab-17246  |
|---------------------------|--|
| Isotype                   | IgG  |
| Reactivity                | Human, Mouse,Rat   |
| Applications              | IHC,WB   |
| Gene Name                 | HMGCL  |
| Protein Name              | Hydroxymethylglutaryl-CoA lyase, mitochondrial (HL) (HMG-CoA lyase) (EC 4.1.3.4) (3-hydroxy-3-methylglutarate-CoA lyase)   |
| Immunogen                 | Synthesized peptide derived from human C-ternal HMGCL  |
| Specificity               | This antibody detects endogenous levels of HMGCL at Human, Mouse,Rat   |
| Formulation               | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| Source                    | Rabbit,polyclonal  |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution                  | WB 1:500-2000 IHC 1:50-200   |
| Concentration             | 1 mg/ml  |
| Purity                    | ≥90%   |
| Storage Stability         | -20°C/1 year   |
| Synonyms                  | Hydroxymethylglutaryl-CoA lyase, mitochondrial (HL) (HMG-CoA lyase) (EC 4.1.3.4) (3-hydroxy-3-methylglutarate-CoA lyase)   |
| Observed Band             |  |
| Cell Pathway              | Mitochondrion matrix . Peroxisome . Unprocessed form is peroxisomal  |
| Tissue Specificity        | Highest expression in liver. Expressed in pancreas, kidney, intestine, testis, fibroblasts and lymphoblasts. Very low expression in brain and skeletal muscle. The relative expression of isoform 2 (at mRNA level) is highest in heart (30%), skeletal muscle (22%), and brain (14%).   |
| Function                  | Mitochondrial 3-hydroxymethyl-3-methylglutaryl-CoA lyase that catalyzes a cation-dependent cleavage of (S)-3-hydroxy-3-methylglutaryl-CoA into acetyl-CoA and acetoacetate, a key step in ketogenesis. Terminal step in leucine catabolism. Ketone bodies (beta-hydroxybutyrate, acetoacetate and acetone) are essential as an alternative source of energy to glucose, as lipid precursors and as regulators of metabolism. |
| Background                |  |
| matters needing attention | Avoid repeated freezing and thawing!   |
|                           |  |



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

| Products Images |
|-----------------|
|                 |
|                 |
|                 |
|                 |