







## TBC16 rabbit pAb


Catalog No	YP-Ab-08208
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TBC1D16
Protein Name	TBC16
Immunogen	Synthesized peptide derived from human TBC16 AA range: 529-579
Specificity	This antibody detects endogenous levels of TBC16 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.323% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year TBC1 domain family member 16
	·
Synonyms	TBC1 domain family member 16
Synonyms Observed Band	TBC1 domain family member 16 85kD
Synonyms Observed Band Cell Pathway	TBC1 domain family member 16 85kD early endosome,cytosol,
Synonyms  Observed Band  Cell Pathway  Tissue Specificity	TBC1 domain family member 16  85kD  early endosome,cytosol,  Brain,Breast,Skin,Urinary bladder,  function:May act as a GTPase-activating protein for Rab family
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function	TBC1 domain family member 16  85kD  early endosome,cytosol,  Brain,Breast,Skin,Urinary bladder,  function:May act as a GTPase-activating protein for Rab family
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function  Background  matters needing	TBC1 domain family member 16  85kD  early endosome,cytosol,  Brain,Breast,Skin,Urinary bladder,  function:May act as a GTPase-activating protein for Rab family protein(s).,similarity:Contains 1 Rab-GAP TBC domain.,







## **Products Images**



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night