







Bek Monoclonal Antibody

Catalog No	YP-mAb-13157
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC-p;IF/ICC;ELISA
Gene Name	FGFR2
Protein Name	Fibroblast growth factor receptor 2
Immunogen	The antiserum was produced against synthesized peptide derived from human FGFR2. AA range:471-520
Specificity	Bek Monoclonal Antibody detects endogenous levels of Bek protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGFR2; BEK; KGFR; KSAM; Fibroblast growth factor receptor 2; FGFR-2; K-sam; KGFR; Keratinocyte growth factor receptor; CD antigen CD332
Observed Band	120
Cell Pathway	Cell membrane; Single-pass type I membrane protein. Golgi apparatus. Cytoplasmic vesicle. Detected on osteoblast plasma membrane lipid rafts. After ligand binding, the activated receptor is rapidly internalized and degraded.; [Isoform 1]: Cell membrane; Single-pass type I membrane protein. After ligand binding, the activated receptor is rapidly internalized and degraded.; [Isoform 3]: Cell membrane; Single-pass type I membrane protein. After ligand binding, the activated receptor is rapidly internalized and degraded.; [Isoform 8]: Secreted.; [Isoform 13]: Secreted.
Tissue Specificity	Blood,Brain,Cerebellum,Cornea,Mammary gland,Neonatal brain stem,Pla
Function	
Background	



UpingBio technology Co.,Ltd

© Tel: 400-999-8863 ■ Email:UpingBio@163.com



matters	needing	
attention		

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images