



E2F-1 mouse mAb

Catalog No	YP-Ab-01106
Isotype	IgG
Reactivity	Human;Rat
Applications	WB;IF;IP
Gene Name	e2f1
Protein Name	
Immunogen	Purified recombinant human E2F-1 protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of E2F-1 and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb dilution 1:500 icc dilution 1:100. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Dmel\CG6376 ; Dmel_CG6376 ; drosE2F1 ; E(Sev-CycE)3A ; E(var)3-93E ; E2-promoter binding facto ; E2F 1 ; E2F transcription factor 1 ; E2F-1 ; E2f-PA ; E2f-PB ; E2f-PC ; E2F1 ; E2f1 E2F transcription factor 1 ; E2F1_HUMAN ; Evar(3)164 ; KIAA4009 ; I(3)07172 ; I(3)j3B1 ; I(3)j3C2 ; I(3)rM729 ; mKIAA4009 ; OTTHUMP00000030661 ; PBR3 ; PRB binding protein E2F 1 ; PRB-binding protein E2F-1 ; RBAP 1 ; RBAP-1 ; RBAP1 ; RBBP-3 ; RBBP3 ; RBP 3 ; RBP3 ; Retinoblastoma-associated protein 1 ; Retinoblastoma-binding protein 3 ; Transcription factor E2F1.
Observed Band	60kD
Cell Pathway	Nucleus .
Tissue Specificity	Brain,Epithelium,Pancreas,Skin,
Function	function:Transcription activator that binds DNA cooperatively with dp proteins through the E2 recognition site, 5'-TTTC[CG]CGC-3' found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DRTF1/E2F complex functions in the control of cell-cycle progression from G1 to S phase. E2F-1 binds preferentially RB1 protein, in a cell-cycle dependent manner. It can mediate both cell proliferation and



p53-dependent apoptosis.,PTM:Phosphorylated by CDK2 and cyclin A-CDK2 in the S-phase.,similarity:Belongs to the E2F/DP family.,subunit:Component of the DRTF1/E2F transcription factor complex. Forms heterodimers with DP family members. The E2F-1 complex binds specifically hypophosphorylated retinoblastoma protein RB1. During the cell cycle, RB1 becomes phosphorylated in mid-to-late G1 phase, detaches from the DRTF1/E2F complex, ren

Background

The protein encoded by this gene is a member of the E2F family of transcription factors. The E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionally conserved domains found in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein and another 2 members, E2F2 and E2F3, have an additional cyclin binding domain. This protein binds preferentially to retinoblastoma protein pRB in a cell-cycle dependent manner. It can media

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western blot detection of E2F-1 in C6,Raw264.7 and HeLa cell lysates using E2F-1 mouse mAb (1:500 diluted).Predicted band size:70KDa.Observed band size:70KDa.



Immunofluorescent analysis of HeLa cells fixed with 4% Paraformaldehyde and using anti-E2F-1 mouse mAb (dilution 1:100). DAPI was used to stain nucleus(blue).



Immunoprecipitation analysis of HeLa cell lysates using E2F-1 mouse mAb.