



AMACR(C-term) mouse mAb

| | |
|---------------------------|--|
| Catalog No | YP-Ab-03457 |
| Isotype | IgG |
| Reactivity | Rat |
| Applications | WB;ICC |
| Gene Name | amacr |
| Protein Name | |
| Immunogen | Purified recombinant human AMACR(C-terminus) protein fragments expressed in E.coli. |
| Specificity | This antibody detects endogenous levels of AMACR(C-terminus) and does not cross-react with related proteins. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen. |
| Dilution | wb 1:1000 icc 1:100 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | 2 arylpropionyl CoA epimerase;2 methylacyl CoA racemase;2-methylacyl-CoA racemase;Alpha methylacyl CoA racemase;Alpha methylacyl Coenzyme A racemase;Alpha methylacyl-CoA racemase deficiency, included;Alpha-methylacyl-CoA racemase;Amacr;AMACR deficiency, included;AMACR_HUMAN;CBAS4;Da1-8;EC 5.1.99.4;Macr 1;Macr1;Methylacyl CoA racemase alpha;P504S;RACE;RM. |
| Observed Band | 42kD |
| Cell Pathway | Peroxisome . Mitochondrion . |
| Tissue Specificity | Aorta,Brain,Cerebellum,Kidney,Liver,PCR rescued clones,Prostate cancer,Sali |
| Function | catalytic activity:(2S)-2-methylacyl-CoA = (2R)-2-methylacyl-CoA.,disease:Defects in AMACR are the cause of alpha-methylacyl-CoA racemase deficiency (AMACRD) [MIM:604489]. AMACRD results in elevated plasma concentrations of pristanic acid C27-bile-acid intermediates. It can be associated with polyneuropathy, retinitis pigmentosa, epilepsy.,disease:Defects in AMACR are the cause of congenital bile acid synthesis defect type 4 (CBAS4) [MIM:214950]; also known as cholestasis, intrahepatic, with defective conversion of trihydroxycoprostanic acid to cholic acid |



or trihydroxycoprostanic acid in bile. Clinical features include neonatal jaundice, intrahepatic cholestasis, bile duct deficiency and absence of cholic acid from bile. function: Racemization of 2-methyl-branched fatty acid CoA esters. Responsible for the conversion of pristanoyl-CoA and C27-bile acyl-CoAs to their (S)-stereoisomers. pa

Background

This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene. [provided by RefSeq, Mar 2011],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot detection of AMACR(C-terminus) in Rat Liver lysates using AMACR(C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 42KDa. Observed band size: 42KDa.



Immunocytochemistry staining of Jurkat cells fixed with -20°C Ethanol and using anti-AMACR (C-terminus) mouse mAb (dilution 1:100).