




# RACK1 mouse mAb

货号	YP-mAb-18321
同位型	IgG
应用	WB
种属	Human, Mouse,Rat
其他名称	H12.3; HLC-7; PIG21; GNB2L1; Gnb2-rs1; RACK1
免疫原	Recombinant fusion protein containing a sequence corresponding to amino acids 220-310 of human RACK1 (P63244)
稀释	WB 1:1000 - 1:6000
纯化工艺	Affinity purification
分子量	35kDa
背景	Enables several functions, including cyclin binding activity; enzyme binding activity; and protein domain specific binding activity. Involved in several processes, including positive regulation of hydrolase activity; regulation of cellular protein metabolic process; and regulation of signal transduction. Located in several cellular components, including midbody; perinuclear region of cytoplasm; and phagocytic cup. Part of IRE1-RACK1-PP2A complex
细胞定位	
组织表达	
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western blot analysis of various lysates using RACK1 Mouse mAb (A3808) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Western blot analysis of various lysates using RACK1 Mouse mAb (A3808) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 $\mu$ g per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.