



SUCLG2 mouse mAb

Catalog No	YP-mAb-18386
Isotype	IgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	SUCLG2
Protein Name	Succinyl-CoA ligase [GDP-forming] subunit beta, mitochondrial (GTP-specific succinyl-CoA synthetase subunit beta) (Succinyl-CoA synthetase beta-G chain) (SCS-betaG)
Immunogen	Synthesized peptide derived from human SUCLG2
Specificity	This antibody detects endogenous levels of SUCLG2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	48kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Mainly expressed in liver, kidney, heart, spleen and skeletal muscle. Also found in intestine and colon, and in low amounts in lung, brain, prostate, testis and ovary.
Function	GTP-specific succinyl-CoA synthetase functions in the citric acid cycle (TCA), coupling the hydrolysis of succinyl-CoA to the synthesis of GTP and thus represents the only step of substrate-level phosphorylation in the TCA. The beta subunit provides nucleotide specificity of the enzyme and binds the substrate succinate, while the binding sites for coenzyme A and phosphate are found in the alpha subunit.
Background	
matters needing attention	Avoid repeated freezing and thawing!



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images