

🔇 Tel: 400-999-8863 💌 Email:3631691544@qq.com

Website: www.upingBio.com

CBLL1 mouse mAb

Catalog No	YP-mAb-18398
Isotype	lgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	CBLL1 HAKAI RNF188
Protein Name	E3 ubiquitin-protein ligase Hakai (Casitas B-lineage lymphoma-transforming sequence-like protein 1) (RING finger protein 188) (c-Cbl-like protein 1)
Immunogen	Synthesized peptide derived from human CBLL1
Specificity	This antibody detects endogenous levels of CBLL1 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year 54kD Nucleus speckle . Nucleus, nucleoplasm . Cytoplasm . Mainly nuclear with some
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 54kD Nucleus speckle . Nucleus, nucleoplasm . Cytoplasm . Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 54kD Nucleus speckle . Nucleus, nucleoplasm . Cytoplasm . Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 ≥90% -20°C/1 year 54kD Nucleus speckle . Nucleus, nucleoplasm . Cytoplasm . Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the MACOM subcomplex, such as VIRMA, in the nucleus E3 ubiquitin-protein ligase that mediates ubiquitination of several tyrosine-phosphorylated Src substrates, including CDH1, CTTN and DOK1 (By similarity). Targets CDH1 for endocytosis and degradation (By similarity). Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing . Its function in the WMM



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 💌 Email:3631691544@qq.com 🤇

WebsIte: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images