

🔇 Tel: 400-999-8863 📼 Emall:3631691544@qq.com 🛛 🙆 Website: www.upingBio.com

## Notch4 Mouse mAb

Catalog No	YP-mAb-18442
lsotype	lgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	
Protein Name	
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1824-2003 of human NOTCH4 (NP_004548.3
Specificity	
Formulation	
Source	
Purification	Affinity purification
Dilution	WB 1:100 - 1:500
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	INT3; NOTCH4
Observed Band	80kDa
Cell Pathway	
Tissue Specificity	
Function	
Background	This gene encodes a member of the NOTCH family of proteins. Members of this Type I transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple different domain types. Notch signaling is an evolutionarily conserved intercellular signaling pathway that regulates interactions between physically adjacent cells through binding of Notch family receptors to their cognate ligands. The encoded preproprotein is proteolytically processed in the trans-Golgi network to generate two polypeptide chains that heterodimerize to form the mature cell-surface receptor. This receptor may play a role in vascular, renal and hepatic development. Mutations in this gene may be associated with schizophrenia. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.



🔇 Tel: 400-999-8863 📼 Email:3631691544@qq.com 🛛 🙆 Wel

Website: www.upingBio.com


matters needing attention

Usage suggestions

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of Mouse lung, using NOTCH4 Mouse pAb (A8303) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.