





TBRG4 Rabbit pAb

Catalog No	YP-Ab-18467
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	TBRG4 CPR2 FASTKD4 KIAA0948
Protein Name	Protein TBRG4 (Cell cycle progression restoration protein 2) (Cell cycle progression protein 2) (FAST kinase domain-containing protein 4) (Transforming growth factor beta regulator 4)
Immunogen	Synthesized peptide derived from human TBRG4
Specificity	This antibody detects endogenous levels of TBRG4 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration	i mg/m
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 69kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 69kD Mitochondrion matrix . Ubiquitously expressed (PubMed:12759187). Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in smooth muscle
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 69kD Mitochondrion matrix . Ubiquitously expressed (PubMed:12759187). Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in smooth muscle (PubMed:20869947). Plays a role in processing of mitochondrial RNA precursors and in stabilization of a subset of mature mitochondrial RNA species, such as MT-CO1, MT-CO2, MT-CYB, MT-CO3, MT-ND3, MT-ND5 and MT-ATP8/6. May play a role in cell
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 69kD Mitochondrion matrix . Ubiquitously expressed (PubMed:12759187). Expression detected in spleen, thymus, testis, ovary, colon, heart, smooth muscle, kidney, brain, lung, liver and white adipose tissue with highest expression in smooth muscle (PubMed:20869947). Plays a role in processing of mitochondrial RNA precursors and in stabilization of a subset of mature mitochondrial RNA species, such as MT-CO1, MT-CO2, MT-CYB, MT-CO3, MT-ND3, MT-ND5 and MT-ATP8/6. May play a role in cell



UpingBio technology Co.,Ltd



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images