

C Tel: 400-999-8863 ■ Email:Upingbio.163.com



## PSA (ABT114) Mouse mAb

| 货号   | YP-Ab-17762   |
|------|---|
| 同位型  | IgG   |
| 靶点   | PSA   |
| 简介   | >>Pathways in cancer;>>Prostate cancer  |
| 基因名称 | KLK3 APS  |
| 种属   | Human;Mouse;Rat   |
| 应用   | IHC;WB;IF   |
| 蛋白名称 | PSA   |
| 免疫原  | Synthesized peptide derived from human PSA AA range: 100-200  |
| 特异性  | The antibody can specifically recognize human PSA protein The antibody was also Predict react with Mouse;Rat  |
| 组成   | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| 来源   | Mouse, Monoclonal/IgG2b, Kappa  |
| 稀释   | IHC-p 1:200-400, WB 1:200-1000, IF 1:200-400  |
| 纯化工艺 | The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.  |
| 其他名称 | Prostate-specific antigen (PSA;EC 3.4.21.77;Gamma-seminoprotein;Seminin;Kallikrein-3;P-30 antigen;Semenogelase)   |
| 背景   | Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Its protein product is a protease present in seminal plasma. It is thought to function normally in the liquefaction of seminal coagulum, presumably by hydrolysis of the high molecular mass seminal vesicle protein. Serum level of this protein, called PSA in the clinical setting, is useful in the diagnosis and monitoring of prostatic carcinoma. Alternate splicing of this gene generates several transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008], |
| 功能   | catalytic activity:Preferential cleavage: -Tyr- -Xaa,function:Hydrolyzes semenogelin-1 thus leading to the liquefaction of the seminal coagulum.,online information:Prostate-specific antigen entry,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family. Kallikrein subfamily.,similarity:Contains 1 peptidase S1 domain.,   |
| 细胞定位 | Secreted.   |
| 组织表达 | Cytoplasmic   |
| 浓度   | 1 mg/ml   |
|      |   |




## UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:Upingbio.163.com



| 储存     | -15°C to -25°C/1 year(Do not lower than -25°C)  |
|--------|---|
| 有关注意事项 | Avoid repeated freezing and thawing!  |
| 使用建议   | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

## **Products Images**







## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com



