



# c-Myc Tag Mouse mAb

<b>Catalog No</b>	YP-Ab-17868
<b>Isotype</b>	IgG1
<b>Reactivity</b>	Species-independent
<b>Applications</b>	WB, ICC/IF, IP
<b>Gene Name</b>	-
<b>Alternative Names</b>	c-Myc; Myc tag; Myc protein; Myc epitope tag; Myc2
<b>Research Field</b>	Epigenetics and Nuclear Signaling
<b>Product Categories</b>	Tag antibody
<b>Host</b>	Mouse
<b>Molecular Weight</b>	Refer to figures
<b>Clonality</b>	Monoclonal Antibody
<b>Clonality No.</b>	7D9-4C6-6F10
<b>Dilution</b>	WB: 1/1000-1/10000 IF: 1/100-1/500 IP: 1/200
<b>Immunogen</b>	Synthetic Peptide of Myc-Tag
<b>Purification</b>	Affinity Purified
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Form</b>	Liquid
<b>Buffer System</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Background</b>	Myc tag is a polypeptide protein tag derived from the c-myc gene product that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, and then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



