




# Rabbit IgG mouse mAb


<b>Catalog No</b>	YP-Ab-04489
<b>Isotype</b>	IgG
<b>Reactivity</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	rabbit igg
<b>Protein Name</b>	
<b>Immunogen</b>	Rabbit IgG
<b>Specificity</b>	This antibody detects rabbit IgG.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	wb dilution 1:50000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Antibody; Ig gamma chain C region; Ig kappa-b4 chain C region; Ig kappa-b5 chain C region; Ig kappa-b9 chain C region; Ig lambda chain C region; K-BAS.
<b>Observed Band</b>	
<b>Cell Pathway</b>	
<b>Tissue Specificity</b>	
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images



Western blot detection of MEK-2 in THP-1, Jurkat, Raji, 3T3 and Hela cell lysates using 310001 (MEK-2 rabbit pAb, 1:1000 diluted), then incubated with 201069 (Rabbit IgG mAb, 1:50000 diluted). Anti-mouse IgG, HRP-linked Antibody was used for detecting. Predicted band size: 44KDa. Observed band size: 44KDa.



Western blot analysis of decreasing concentrations of total rabbit IgG reduced and denatured in 2X SDS loading buffer with DTT using Mouse Anti-Rabbit IgG (Heavy-Chain Specific) mAb (1:1000 diluted).



Immunoprecipitation analysis of CHO-K1 cell lysates transfected with HA-tagged NGAL using HA-Tag Rabbit pAb (390001, 1:100 diluted) (lane 2). Lane 1 contains rabbit IgG IP control lysate. All lanes are probed with HA-Tag Rabbit pAb (390001, 1:1000 diluted) as the primary antibody. Panel A incubated with secondary antibodies Anti-Rabbit whole IgG, HRP-linked Antibody for detecting. Panel B incubated with Anti-Rabbit IgG (Heavy-Chain Specific) mAb (1:50000 diluted) and the Anti-mouse whole IgG, HRP-linked Antibody was used for detecting.