




HptII mouse mAb

| | |
|----------------------------------|---|
| Catalog No | YP-Ab-04490 |
| Isotype | IgG |
| Reactivity | Transgenic |
| Applications | WB |
| Gene Name | hptii |
| Protein Name | |
| Immunogen | |
| Specificity | Transgenic Only. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse |
| Purification | The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen. |
| Dilution | wb dilution 1:1000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 37kD |
| Cell Pathway | |
| Tissue Specificity | |
| Function | |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |



Products Images



Western blot detection of HptII in Transgenic rice (leaf) and Non-transgenic rice (leaf) cell lysates using HptII mouse mAb (1:1000 diluted). Predicted band size: 36,38KDa. Observed band size: 36,38KDa. Beta-actin mouse mAb(200068-8F10, 1:1000 diluted) was used as a loading control.