



PCSK9 Mouse mAb(5G4)

Catalog No	YP-Ab-04835
Isotype	IgG
Reactivity	Human
Applications	IHC
Gene Name	PCSK9 NARC1 PSEC0052
Protein Name	PCSK9
Immunogen	Synthesized peptide derived from human PCSK9
Specificity	This antibody detects endogenous levels of PCSK9 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.84% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Proprotein convertase subtilisin/kexin type 9 (EC 3.4.21.-) (Neural apoptosis-regulated convertase 1) (NARC-1) (Proprotein convertase 9) (PC9) (Subtilisin/kexin-like protease PC9)
Observed Band	74kD
Cell Pathway	Cytoplasm. Secreted. Endosome. Lysosome. Cell surface. Endoplasmic reticulum. Golgi apparatus. Autocatalytic cleavage is required to transport it from the endoplasmic reticulum to the Golgi apparatus and for the secretion of the mature protein. Localizes to the endoplasmic reticulum in the absence of LDLR and colocalizes to the cell surface and to the endosomes/lysosomes in the presence of LDLR. The sorting to the cell surface and endosomes is required in order to fully promote LDLR degradation.
Tissue Specificity	Expressed in neuro-epithelioma, colon carcinoma, hepatic and pancreatic cell lines, and in Schwann cells.
Function	cofactor:Calcium.,disease:Defects in PCSK9 are the cause of familial hypercholesterolemia 3 (FH3) [MIM:603776]. FH3 inheritance is autosomal dominant.,enzyme regulation:Inhibited by EGTA.,function:May be implicated in the differentiation of cortical neurons and may play a role in cholesterol homeostasis.,PTM:The soluble zymogen undergoes autocatalytic intramolecular processing in the endoplasmic reticulum, resulting in the cleavage of its propeptide that remains associated with the secreted enzyme.,similarity:Belongs to the peptidase S8 family.,similarity:Contains 1 peptidase S8



domain.,subunit:The precursor protein but not the mature protein may form multimers.,tissue specificity:Expressed in neuro-epithelioma, colon carcinoma, hepatic and pancreatic cell lines, and in Schwann cells.,

Background

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver, intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded Human Kidney Tissue using PSCK9 Mouse Monoclonal antibody diluted at 1:200.